

DePauw University

## Scholarly and Creative Work from DePauw University

---

Science Research Fellows Posters

Student Work

---

Fall 10-2020

### Analysis of Dynamical Systems for Synthesis of Phenobarbital

Mishal Ali  
*DePauw University*

Follow this and additional works at: <https://scholarship.depauw.edu/srfposters>



Part of the [Applied Mathematics Commons](#), and the [Chemistry Commons](#)

---

#### Recommended Citation

Ali, Mishal, "Analysis of Dynamical Systems for Synthesis of Phenobarbital" (2020). *Science Research Fellows Posters*. 42.

<https://scholarship.depauw.edu/srfposters/42>

This Poster is brought to you for free and open access by the Student Work at Scholarly and Creative Work from DePauw University. It has been accepted for inclusion in Science Research Fellows Posters by an authorized administrator of Scholarly and Creative Work from DePauw University. For more information, please contact [bcox@depauw.edu](mailto:bcox@depauw.edu).



## OBJECTIVES


1. Understanding sequential chemical reactions.
2. Understanding basic mathematical theories in the Linear Algebra and Differential Equations.
3. Investigating the synthesis of industrial aromatic compounds and experiencing their visualization using computational programs such as Python and Mathematica.
4. Enhancing the ability to analyze the phenomena in the chemical reactions using the data coming from the mathematical works.

## INTRODUCTION

The use of mathematical methods for the analysis of chemical reaction systems is one of the useful tools. Phenobarbital (a barbiturate type medication also called phenobarb) is a prescription drug used to control seizures, relieve anxiety, treat epilepsy (in some countries), and prevent withdrawal symptoms in people dependent on other barbiturate drugs. We followed the strategy employed in the paper [1] but approaches it with different mathematical approaches: matrix analysis [2] and ODE system [3] which will help us understand the chemical stoichiometry of these synthesis reactions.

## CHEMICAL MODEL

We consider the following sequential chemical reactions showing the synthesis of Phenobarbital as the final product



## MATHEMATICAL MODEL

Also, the following system of ordinary differential equation (ODE System) led to the deduction:

$$\begin{aligned}\frac{dA_0}{dt} &= -k[A_0] \\ \frac{dA_1}{dt} &= k[A_0] - k[A_1] \\ \frac{dA_2}{dt} &= k[A_1] - k[A_2] \\ \frac{dA_3}{dt} &= k[A_2] - k[A_3] \\ \frac{dA_4}{dt} &= -k[A_4]\end{aligned}$$

## MATHEMATICAL METHODS

The main idea to solve the ODE System is to consider the following matrix:

$$\frac{d\vec{x}}{dt}(t) = A\vec{x}(t)$$

where

$$\vec{x}(t) = \begin{pmatrix} A_0(t) \\ A_1(t) \\ A_2(t) \\ A_3(t) \\ A_4(t) \end{pmatrix},$$

$$A = \begin{pmatrix} -k & 0 & 0 & 0 & 0 \\ k & -k & 0 & 0 & 0 \\ 0 & k & -k & 0 & 0 \\ 0 & 0 & k & -k & 0 \\ 0 & 0 & 0 & 0 & k \end{pmatrix}.$$

## MATHEMATICAL AND NUMERICAL RESULT

Using principles and techniques from Linear Algebra we are able to find the eigenvalues and its corresponding eigenvectors for the matrix which are key elements to finding the general solution.

$$\lambda_1 = 0, \lambda_2 = -k, \text{ (algebraic multiplicity } = 4), \quad (1)$$

$$\vec{x}_1 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}, \vec{x}_2 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \\ -1 \end{pmatrix}, \vec{x}_3 = \begin{pmatrix} 0 \\ 0 \\ 1/k \\ -1/2k \\ -1/2k \end{pmatrix},$$

$$\vec{x}_4 = \begin{pmatrix} 0 \\ 1/k^2 \\ -1/2k^2 \\ -1/4k^2 \\ -1/4k^2 \end{pmatrix}, \vec{x}_5 = \begin{pmatrix} 1/k^2 \\ -1/2k^2 \\ -1/4k^2 \\ -1/8k^2 \\ -1/8k^2 \end{pmatrix}$$

Finally, the eigenvalues and eigenvectors give us the general solution of the ODE system:

$$\begin{aligned}A_0(t) &= c_1 [1/k^2] e^{-kt} \\ A_1(t) &= c_1 [-1/2k^2] e^{-kt} + c_2 [-1/k^2] t e^{-kt} \\ A_2(t) &= c_1 [-1/4k^2] e^{-kt} + c_2 [-1/2k^2] t e^{-kt} \\ &\quad + c_3 [-1/k] [t^2 e^{-kt/2}] \\ A_3(t) &= c_1 [-1/8k^2] e^{-kt} + c_2 [-1/4k^2] t e^{-kt} \\ &\quad + c_3 [-1/2k] [t^2 e^{-kt/2}] + c_4 [t^3 e^{-kt}] / 6 \\ A_4(t) &= c_1 [-1/8k^2] e^{-kt} + c_2 [-1/4k^2] t e^{-kt} \\ &\quad + c_3 [-1/2k] [t^2 e^{-kt/2}] + c_4 [t^3 e^{-kt}] / 6.\end{aligned}$$

## REFERENCES

- [1] Jack McGeachy. The progression of sequential reactions. *Undergraduate Journal of Mathematical Modeling: One + Two*, 2:Iss 2, Article 5, 2010.
- [2] Iljas Farah and Ken Kuttler. *A First Course in Linear Algebra*. Ilyx Learning, free edition edition, 2017.
- [3] William E. Trench. *Elementary Differential Equations with Boundary Value Problem*. Brooks/Cole Thomson Learning, free edition edition, 2013.

## NUMERICAL SIMULATION AND CONCLUSION




Figure 1: The Shape of Solution Graphs as  $k = 0.5$

- The mathematical works and numerical visualization allow us find a Maximum Points for each solution function and intermediate involved in the reaction.
- Our analysis shows that chemical stoichiometry of Phenobarbital can be approximated from these results.

## ACKNOWLEDGEMENT

- This project has been advised by Professor Seonguk Kim, Assistant Professor of Mathematics.
- The project was supported by the Science Research Fellows Program at DePauw University and performed for summer research project 2020.